

# Dietary glycemic load in adolescent girls with and without acne

Safoura Shakoei, MD <sup>1</sup>  
Nazanin Nasri, MD <sup>2</sup>  
Ghazal Shariatpanahi, MD <sup>2</sup>  
Rezvan Hashemi, MD, PhD <sup>3\*</sup>

1. Department of Dermatology, Imam Khomeini Hospital, Tehran University of Medical Sciences, Tehran, Iran
2. Department of Pediatrics, Ziaeiian Hospital, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran
3. Department of Geriatric Medicine, Ziaeiian Hospital, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

\*Corresponding author:  
Rezvan Hashemi, MD  
Department of Geriatric Medicine,  
Ziaeiian Hospital, School of Medicine,  
Tehran University of Medical Sciences,  
Tehran, Iran  
Tel: +982155176810  
Fax: +982155176007  
E-mail: rezvan.hme@gmail.com

Received: 22 January 2020  
Accepted: 26 February 2020

## INTRODUCTION

Acne is a common inflammatory skin disease, which mostly presents during adolescence. It affects a significant proportion of adolescents, with a prevalence of 79-95% in Western countries <sup>1,2</sup>. The prevalence of acne has increased among adults in developed countries. It may present, persist, or recur in the third and fourth decades of life <sup>3</sup>.

It predominantly affects the skin and is consequently associated with psychological distress, low self-esteem, poor body image, social

**Background:** Acne is a chronic inflammatory skin disease and a cosmetic problem with considerable emotional and psychological side effects and symptoms, such as pain and pruritus. Some controversies exist concerning the involvement of dietary factors, including glycemic load (GL), in the pathogenesis of acne. Accordingly, we aimed to determine the role of GL and anthropometric measures in acne development among girls aged 12-18 years in Tehran, Iran.

**Methods:** In this cross-sectional study, 99 girls (45 girls without acne and 54 girls with acne) aged 12-18 years completed a three-day food record. Anthropometric measures, such as height, weight, waist circumference, and body mass index (BMI), were also assessed. Additionally, an expert dermatologist scored the severity of acne, and then, anthropometric measures and GL were examined in the groups.

**Results:** The results showed no significant difference in dietary GL, height, weight, BMI, and waist circumference between the groups. Furthermore, the severity of acne was not significantly associated with GL.

**Conclusion:** The present results did not confirm the association between acne and dietary carbohydrates, including GL. However, further research can contribute to determination of the effect of diet on acne and its severity.

**Keywords:** acne vulgaris, diet, glycemic load, anthropometry

Iran J Dermatol 2020; 23: 43-48

withdrawal, and depression <sup>4,5</sup>. The pathogenesis of acne is complex and attributed to excessive sebum production, follicular hyperkeratinization, intra-follicular colonization by bacteria and yeast, and inflammation <sup>3,6</sup>.

There is some evidence regarding the involvement of environmental factors, including diet, in the acne pathogenesis <sup>6</sup>. Various hormonal mediators, such as insulin-like growth factor-1 (IGF-1), insulin-like growth factor binding protein-3 (IGFBP-3), and sex hormone-binding globulin (SHBG), are involved in acne development by promoting the growth of

keratinocytes and sebaceous glands, along with sebum production<sup>7,8</sup>. There is also controversial evidence regarding the association of dietary patterns with acne pathogenesis. Some studies have indicated the impact of dietary glycemic load (GL) on acne<sup>9</sup>, while some have reported different results<sup>10,11</sup>. Therefore, the role of GL is particularly intriguing and requires major consideration<sup>8</sup>.

Glycemic load refers to the type and amount of consumed carbohydrates and is interpreted as an index of diet-induced insulin demand<sup>9</sup>. A low glycemic index is associated with the reduction of insulin resistance. Therefore, GL can be implicated in acne development due to diet-induced hyperinsulinemia, in addition to the subsequent cascade of hormonal responses and acne exacerbation. Insulin resistance and increased IGF-1 levels change the concentrations of circulating hormones, binding proteins, and receptors, which are associated with the stimulation of unregulated tissue growth androgens, sebum production, and acne-promoting pathways<sup>9</sup>.

The role of dietary carbohydrates in acne pathogenesis has not been established yet. Evidence suggests no significant difference in the level of glucose, insulin, and dietary GL between acne patients and healthy control individuals<sup>10,12</sup>. Considering the high prevalence of acne during adolescence and its negative effects on self-image and self-esteem, it is necessary to determine the effective factors in acne, including dietary GL. In this study, we aimed to investigate the possible association of dietary GL with acne in 12- to 18-year-old girls, using a validated questionnaire for GL.

## PARTICIPANTS AND METHODS

### Participants and study design

Fifty-four females with acne, aged 12-18 years, were recruited in this study. Forty-five age-matched controls were also selected from the Dermatology Outpatient Department of Ziaei Hospital, which is one of the educational centers affiliated with Tehran University of Medical Sciences, Tehran, Iran. Two groups of patients were excluded from the study: acne patients with a history of acne treatment and patients with a history of systemic disorders, such as diabetes. Patients with acne were divided into three groups by an expert dermatologist: mild

(comedonal acne), moderate (papular and pustular lesions), and severe (nodular lesions).

### Anthropometric and clinical data collection

Demographic and anthropometric measures, including age, waist circumference, weight, body mass index (BMI), and height, were recorded for all participants. Height (cm) was measured using a wall tape in a standing position without shoes. Weight (kg) was measured using a digital scale with minimal clothing. Furthermore, waist circumference was measured at the mid-point between the lower rib margin and the iliac crest in a standing position with normal breathing. Moreover, the participants' dietary intake was determined using a three-day food record questionnaire, in which they were asked to document their food, snack, and beverage intake completely over three days (two weekdays and one weekend). An expert nutritionist guided them on how to record their food intake. The questionnaires were completed after one week<sup>13</sup>.

All subjects were informed about the study procedures and objectives and provided an informed consent to participate in the study. Additionally, the participants were ensured about the confidentiality of their information. The study protocol was approved by the Ethics Committee of Tehran University of Medical Sciences and performed in accordance with the Declaration of Helsinki (IR TUMS VCR REC13951849).

The glycemic index of foods was determined, using the International Tables of Glycemic Index and GL Tables<sup>14</sup>. The total carbohydrate and fiber contents of dietary intake were calculated, using the United States Department of Agriculture (USDA) Tables<sup>15</sup>. The dietary information, collected from the food diaries, were converted to grams, and the content of carbohydrate, fiber, and other food components was calculated, using the Modified Nutritionist 4 software. We used the three-day food records to calculate daily GL as follows:

Daily GL = GI for food item × carbohydrate content (g)/100

### Statistical methods

The sample size was calculated with an alpha level of 0.05 and beta level of at least 80% according to previous studies<sup>13</sup>. The calculated sample size

for each group was measured to be 40. Considering the possibility of sample attrition, 55 participants were included in each group. During the study, one patient from the acne group and 10 individuals from the control group withdrew from the study; accordingly, their incomplete information was not included in the final report. Data were analyzed and compared between the two groups, using student's t-test and one-way analysis of variance (ANOVA) in the SPSS software version 24. P value less than 0.05 was considered statistically significant.

## RESULTS

A total of 99 participants were investigated in two groups: acne group (n = 54) and control group (n = 45). Height, weight, waist circumference, and BMI were calculated in the acne and control groups. Our results showed no significant difference between the groups regarding any of the mentioned measurements (Table 1).

Assessment of the adolescents' dietary intake revealed that generally GL ranged from 55.33 to 199.33, with a mean of  $99.91 \pm 25.24$ . The mean GL was not significantly different between the two study groups ( $98.45 \pm 24.88$  vs.  $101.67 \pm 25.85$ ;  $P = 0.903$ ; Table 2). The association between dietary GL and acne severity was examined in the acne group. ANOVA test did not demonstrate any significant differences in the average GL between the three acne groups ( $P = 0.406$ ).

**Table 1.** General and demographic characteristics of the two groups.

Characteristics	Acne (n=54)	Control (n=45)	P-value
	Mean $\pm$ SD		
Age (years)	$15.07 \pm 1.97$	$14.84 \pm 14.84$	0.43
Height (cm)	$159.54 \pm 5.74$	$159.62 \pm 7.26$	0.51
Weight (kg)	$59.85 \pm 13.79$	$57.96 \pm 14.70$	0.94
BMI (kg/m <sup>2</sup> )	$23.37 \pm 4.56$	$22.77 \pm 5.02$	0.77
Waist (cm)	$73.70 \pm 7.04$	$74.18 \pm 10.04$	0.59

BMI, body mass index; cm, centimeter; kg, kilogram; n, number; SD, standard deviation

**Table 2.** Glycemic load calculated in the control and acne groups.

	Control (n=45)	Acne (n=54)	Mild acne	Moderate acne	Severe acne
	Mean $\pm$ SD		Mean $\pm$ SD		
Glycemic load	$101.67 \pm 25.85$	$98.45 \pm 24.88$	$88.98 \pm 16.7$	$101.77 \pm 28.48$	$96.30 \pm 18.70$
P-value	0.903		0.406		

n, number; SD, standard deviation

Only eight patients had mild acne with an average GL of  $88.98 \pm 16.7$ . Thirty-two patients were diagnosed with moderate acne, whose average GL was calculated to be  $101.77 \pm 28.48$ . Finally, 14 patients were diagnosed with severe acne with an average GL of  $96.30 \pm 18.70$  (Table 2).

## DISCUSSION

Our results indicated that dietary GL in Iranian adolescent girls with acne was not higher than that of control subjects. Additionally, no significant association was found between dietary GL and acne severity. Furthermore, BMI and waist circumference of the two groups were not significantly different.

Acne is a common skin disorder affected by different factors, such as genetics<sup>16</sup>, lifestyle and habits<sup>16,17</sup>, seasonal changes<sup>17</sup>, obesity<sup>18,19</sup>, and diet<sup>2,9,20</sup>. However, there are disagreements about the impact of each factor. Many studies support the association between dietary carbohydrates and acne development<sup>9,20-24</sup>. In this regard, a study compared the inhabitants of Kitua Island (Tanzania) and Paraguay hunters, consuming large amounts of vegetable and meat (with lower levels of carbohydrates), with individuals consuming a Western diet; it was found that acne was rarely found in these populations. Therefore, a low-glycemic diet can protect against acne owing to the absence of hyperinsulinemia and activation of the subsequent endocrine cascade<sup>2</sup>.

Ismail *et al.* found an association between dietary GL and acne in Malaysian young adults. Their study showed that GL was significantly higher in the acne group than in the healthy controls, aged 18-30 years. They reported no significant difference in BMI between the case and control groups<sup>13</sup>. Another community-based study reported that individuals maintaining a Mediterranean diet (a low-carbohydrate diet) were less likely to develop acne<sup>25</sup>. Moreover, a cross-sectional study showed that dietary GL was higher among individuals with

moderate to severe acne (mean age =  $21.8 \pm 3.5$  years), compared to those with mild acnes<sup>9</sup>.

The mechanism underlying the association between dietary GL and acne has been well described. A high-GL diet causes hyperinsulinemia with subsequent activation of a signaling cascade, leading to an increase in insulin and IGF-1 activities while reducing IGFBP-3<sup>9,25</sup>. Reduction of IGFBP-3 significantly increases the bioavailability of IGF-1 as a key stimulator in acne pathogenesis<sup>26,27</sup>. IGF-1 stimulates lipogenesis and proliferation of keratinocytes and sebocytes<sup>28,29</sup>. Both insulin and IGF-1 increase the synthesis of gonadal and adrenal androgens<sup>30</sup>. Moreover, IGF-1 reduces the synthesis of SHBG in the liver and occupies androgenic receptors. Consequently, it directly increases the bioavailability of androgens<sup>31,32</sup>. Androgens promote the production of fat as a factor playing a role in acne development. In addition, IGFBP is a strong pro-apoptotic factor in keratinocytes<sup>33</sup>.

Although several studies have been conducted in this area, the association between dietary carbohydrates and acne has not been approved in any investigations, including the current study. In this regard, Kaymak *et al.* in a prospective cohort study, reported no significant difference in terms of serum glucose, insulin, and leptin content, overall glycemic index, and dietary GL between university students with and without acne. It should be noted that they only focused on dietary carbohydrates, similar to our study. However, they suggested a relationship between high glycemic index and duration of acne, which was not addressed in the current study<sup>10</sup>.

Furthermore, the results of a non-randomized clinical trial demonstrated that acne improved in adolescent boys on both low- and high-GL diets, while the difference between the groups was not significant. They found no significant correlation between changes in acne severity and insulin concentration or sensitivity. They also reported that diet significantly affected IGFBP-1, but not IGFBP-3 or IGF-1<sup>11</sup>. In this respect, Smith *et al.* conducted four interventional studies to investigate the effect of dietary GL on acne severity. In two clinical trials, low dietary GL significantly reduced acne severity and plasma levels of free androgens, while it increased insulin and IGFBP sensitivity<sup>34,35</sup>.

Consistent with the findings reported by Smith *et al.*, recent studies including a clinical

trial, have provided histopathological evidence regarding the benefits of a low-GL regimen in reducing acne development. They found that low GL was associated with reduced acne severity, sebum production, expression of sterol regulatory element-binding proteins, lipid synthesis, and inflammatory cytokines like interleukin-8<sup>9</sup>. Nevertheless, Pavithra *et al.* did not find any improvement in acne development following adherence to a low-GL diet<sup>23</sup>. Overall, the impact of weight loss secondary to a low-GL diet has not been examined in any of the mentioned studies, while BMI is an important contributor to acne development.

In a number of studies, the effect of low-GL diet could not be detected when the data were statistically adjusted for BMI changes<sup>13</sup>. In the current study, there was no significant difference in BMI between adolescents with and without acne, which could explain the lack of association between GL and acne development. Additionally, some previous studies demonstrated no significant correlation between BMI and presence of acne or its severity<sup>13, 36-38</sup>.

The Iranian dietary pattern is high in refined grains, including white rice and bread, potato, tea, whole grains, hydrogenated fats, legumes, and broth, and differs from Mediterranean and Western dietary patterns<sup>39</sup>. It has been revealed that the Iranian diet is not associated with general or central obesity<sup>39</sup>, and the mean GL in the present study is similar to that of previous reports. Although many foods rich in carbohydrates are included in the typical Iranian diet, it appears that some other healthy food groups in this diet, such as legumes and whole grains, could alleviate the effect of high-carbohydrate components<sup>39</sup>. Other factors, such as genetics<sup>16</sup>, tobacco use, and different climatic conditions<sup>17</sup> may be also influential in acne development. Furthermore, differences in the age range of participants may explain the discrepancy between our findings and previous reports.

This study had some limitations. First, we used a three-day food record owing to time and cost constraints, while other food recording methods, such as repeated three-day questionnaire and food frequency questionnaire, would be more effective in calculating GL, especially in patients with chronic diseases<sup>13,40</sup>. Second, the sample size was calculated to be sufficient to analyze the differences based

on independent t-test. Consequently, the sample size was not large enough for advanced statistical analyses.

## CONCLUSION

Chronic skin diseases, including acne, are often accompanied by significant emotional and psychological burdens negatively affecting the individual's quality of life. Thus, control of modifiable factors to reduce acne is necessary. Change of dietary habits is a controversial treatment option for patients with acne. The results of the present study did not confirm the effects of dietary GL on acne in young girls. Moreover, no significant difference in BMI and waist circumference was observed between the acne and healthy groups.

## Acknowledgements

The authors sincerely appreciate Ziaieian Hospital for recruiting the participants.

**Conflict of Interest:** None declared.

## REFERENCES

- Ghodsi SZ, Orawa H, Zouboulis CC. Prevalence, severity, and severity risk factors of acne in high school pupils: a community-based study. *J Invest Dermatol.* 2009;129(9):2136-41.
- Cordain L, Lindeberg S, Hurtado M, et al. Acne vulgaris: a disease of Western civilization. *Arch Dermatol.* 2002;138(12):1584-90.
- Williams HC, Dellavalle RP, Garner S. Acne vulgaris. *The Lancet.* 2012; 379(9813):361-72.
- Gieler U, Gieler T, Kupfer J. Acne and quality of life—impact and management. *J Eur Acad Dermatol Venereol.* 2015 (Suppl 4:);29:12-4.
- Tan J. Psychosocial impact of acne vulgaris: evaluating the evidence. *Skin Therapy Lett.* 2004;9(7):1-3,9.
- Burris J, Rietkerk W, Woolf K. Acne: the role of medical nutrition therapy. *J Acad Nutr Diet.* 2013;113(3):416-30.
- Kurokawa I, Danby FW, Ju Q, et al. New developments in our understanding of acne pathogenesis and treatment. *Exp Dermatol.* 2009;18(10):821-32.
- Bowe WP, Joshi SS, Shalita AR. Diet and acne. *J Am Acad Dermatol.* 2010;63(1):124-41.
- Burris J, Rietkerk W, Shikany JM, et al. Differences in dietary glycemic load and hormones in New York city adults with no and moderate/severe acne. *J Acad Nutr Diet.* 2017;117(9):1375-83.
- Kaymak Y, Adisen E, Ilter N, et al. Dietary glycemic index and glucose, insulin, insulin-like growth factor-1, insulin-like growth factor binding protein 3, and leptin levels in patients with acne. *J Am Acad Dermatol.* 2007;57(5):819-23.
- Reynolds RC, Lee S, Choi JY, et al. Effect of the glycemic index of carbohydrates on acne vulgaris. *Nutrients.* 2010;2(10):1060-72.
- Çerman AA, Aktaş E, Altunay İK, et al. Dietary glycemic factors, insulin resistance, and adiponectin levels in acne vulgaris. *J Am Acad Dermatol.* 2016;75(1):155-62.
- Ismail NH, Manaf ZA, Azizan NZ. High glycemic load diet, milk and ice cream consumption are related to acne vulgaris in Malaysian young adults: a case control study. *BMC dermatol.* 2012;12(1):13.
- Atkinson FS, Foster-Powell K, Brand-Miller JC. International tables of glycemic index and glycemic load values: 2008. *Diabetes Care.* 2008;31(12):2281-3.
- US Department of Agriculture, Agricultural Research Service. 2018. USDA food and nutrient database for dietary studies 2015-2016.
- Wolkenstein P, Machovcová A, Szepietowski JC, et al. Acne prevalence and associations with lifestyle: a cross-sectional online survey of adolescents/young adults in 7 European countries. *J Eur Acad Dermatol Venereol.* 2018;32(2):298-306.
- Abo El-Fetoh NM, Alenezi NG, Alshamari NG, et al. Epidemiology of acne vulgaris in adolescent male students in Arar, Kingdom of Saudi Arabia. *J Egypt Public Health Assoc.* 2016; 91(3):144-9.
- Sas K, Reich A. High body mass index is a risk factor for acne severity in adolescents: a preliminary report. *Acta Dermatovenerol Croat.* 2019;27(2):81-5.
- Anaba LE, Ogunbiyi OA, George OA. Adolescent facial acne vulgaris and body mass index: any relationship? *West Afr J Med.* 2019;36(2):129-32.
- Halvorsen JA, Dalgard F, Thoresen M, et al. Is the association between acne and mental distress influenced by diet? results from a cross-sectional population study among 3775 late adolescents in Oslo, Norway. *BMC Public Health.* 2009;9(1):340.
- Law M, Chuh A, Molinari N, et al. An investigation of the association between diet and occurrence of acne: a rational approach from a traditional Chinese medicine perspective. *Clin Exp Dermatol.* 2010;35(1):31-5.
- Rouhani P, Berman B, Rouhani G. Acne improves with a popular, low glycemic diet from South Beach. *J Am Acad Dermatol.* 2009; 60(3):14.
- Pavithra G, Upadya GM, Rukmini MS. A randomized controlled trial of topical benzoyl peroxide 2.5% gel with a low glycemic load diet versus topical benzoyl peroxide 2.5% gel with a normal diet in acne (grades 1-3). *Indian J Dermatol Venereol Leprol.* 2019;85(5):486-90.
- Burris J, Shikany JM, Rietkerk W, et al. A low glycemic index and glycemic load diet decreases insulin-like growth factor-1 among adults with moderate and severe acne: a short-duration, 2-week randomized controlled trial. *J Acad Nutr Diet.* 2018;118(10):1874-85.
- Skroza N, Tolino E, Semyonov L, et al. Mediterranean diet and familial dysmetabolism as factors influencing



- the development of acne. *Scand J Public Health.* 2012;40(5):466-74.
26. Shaw JC. Acne: effect of hormones on pathogenesis and management. *Am J Clin Dermatol.* 2002;3(8):571-8.
27. Vora S, Ovhal A, Jerajani H, et al. Correlation of facial sebum to serum insulin like growth factor-1 in patients with acne. *Br J Dermatol.* 2008;159(4):990-1.
28. Zouboulis CC. Acne vulgaris. *Hautarzt.* 2014;65(8):733-47; quiz 748-9.
29. Thiboutot D. Acne: hormonal concepts and therapy. *Clin Dermatol.* 2004;22(5):419-28.
30. Melnik BC, Schmitz G. Role of insulin, insulin-like growth factor-1, hyperglycaemic food and milk consumption in the pathogenesis of acne vulgaris. *Exp Dermatol.* 2009;18(10):833-41.
31. Berra B, Rizzo AM. Glycemic index, glycemic load: new evidence for a link with acne. *J Am Coll Nutr.* 2009;28 Suppl:450S-454S.
32. Taylor M, Gonzalez M, Porter R. Pathways to inflammation: acne pathophysiology. *Eur J Dermatol.* 2011;21(3):323-33.
33. Cordain L. Implications for the role of diet in acne. *Semin Cutan Med Surg.* 2005;24(2):84-91.
34. Smith RN, Mann NJ, Braue A, et al. A low-glycemic-load diet improves symptoms in acne vulgaris patients: a randomized controlled trial. *Am J Clin Nutr.* 2007;86(1):107-15.
35. Smith RN, Braue A, Varigos GA, et al. The effect of a low glycemic load diet on acne vulgaris and the fatty acid composition of skin surface triglycerides. *J Dermatol Sci.* 2008;50(1):41-52.
36. Hunter N, Shaker O. Diet and body mass index, do they have a role in acne vulgaris? a study using serum leptin and serum Insulin like growth factor-1. *Egypt J Derm Androl.* 2009;9(2):73-80.
37. Lajevardi V, Ghodsi SZ, Daneshpazhooh M, et al. The relationship between body mass index and the severity of acne. *Iran J Dermatol.* 2014;17(1):13-17.
38. Neupane S, Basnet B, Sharma TD. Association between acne and body mass index: a hospital based cross sectional study. *Nepal J Derm Venereol Leprol.* 2018;16(1):53-6.
39. Esmailzadeh A, Azadbakht L. Major dietary patterns in relation to general obesity and central adiposity among Iranian women. *J Nutr.* 2008;138(2):358-63.
40. Basiotis PP, Welsh SO, Cronin FJ, et al. Number of days of food intake records required to estimate individual and group nutrient intakes with defined confidence. *The J Nutr.* 1987;117(9):1638-41.